# Supplementary Files

#### Online Resource 1

Definition (based on ICNARC report on COVID-19 in critical care)(1)

**Comorbidities** must have been evident within the six months prior to critical care and documented at or prior to critical care:

- Cardiovascular: symptoms at rest
- Respiratory: shortness of breath with light activity or home ventilation
- Renal: renal replacement therapy for end-stage renal disease
- Liver: biopsy-proven cirrhosis, portal hypertension or hepatic encephalopathy
- Metastatic disease: distant metastases
- Haematological malignancy: acute or chronic leukaemia, multiple myeloma or lymphoma
- Immunocompromise: chemotherapy, radiotherapy or daily high dose steroid treatment in previous six months, HIV/AIDS or congenital immune deficiency
- Type II diabetes mellitus

**Mechanical ventilation during the first 24 hours** was identified by the recording of a ventilated respiratory rate, indicating that all or some of the breaths or a portion of the breaths (pressure support) were delivered by a mechanical device. This usually indicates invasive ventilation; BPAP (bilevel positive airway pressure) would meet this definition but CPAP (continuous positive airway pressure) does not.

**Advanced respiratory support** was defined as invasive ventilation, BPAP via trans-laryngeal tube or tracheostomy, CPAP via trans-laryngeal tube, extracorporeal respiratory support.

Study site	Viral testing panel
Nottingham University Hospitals	Influenza A & B, RSV, Rhinovirus, Enterovirus, Adenovirus, Parechovirus, Parainfluenza pool (types 1-4), Human metapneumovirus, Bocavirus
Newcastle Upon Tyne Hospitals	Influenza A & B, Respiratory syncytial virus (RSV), Rhinovirus, Human metapneumovirus, Adenovirus, Parainfluenza pool (types 1-4)
Brighton and Sussex University Hospitals	Influenza A & B, RSV
Guy's & St Thomas'	Influenza A & B, RSV, Enterovirus, Rhinovirus, Parainfluenza, Adenovirus, Human metapneumovirus
Salford Royal	Influenza A & B, RSV
University Hospitals of Derby & Burton	Influenza A & B, RSV, Parainfluenza, Rhinovirus, Human metapneumovirus, Adenovirus
University College London	Influenza A & B, RSV, Parainfluenza pool (types 1-4), Human metapneumovirus, Adenovirus, Rhinovirus

Study site	Met inclusion criteria <sup>a</sup>	Still in ICU <sup>b</sup>	Transfers from other hospital-acquired COVID-19°	Eligible <sup>d</sup>	Entered into database (% of those eligible)
Nottingham University Hospitals	97	11	1	85	79 (92.9)
Newcastle Upon Tyne Hospitals	100	4	20	76	48 (63.2)
Brighton and Sussex University Hospitals	58	8	4	46	45 (64.3)
Guy's & St Thomas' NHS Foundation Trust	316	0	103	213	34 (16.0)
Salford Royal	46	3	11	32	22 (68.8)
University Hospitals of Derby & Burton	54	1	0	53	16 (30.2)
University College Hospitals London	138	32	32	74	10 (13.5)
Total	809	59	171	579	254 (43.9)

d= a- (b+c)

Table S 1: Results and classification as likely pathogen or contaminant among positive cultures taken from patients

Blood culture					
Likely pathogen	n	N	Likely contaminant	n	N
Coagulase negative Staphylococcus	13	6	Coagulase negative Staphylococcus	47	36
Enterococcus spp.	8	5	Propionibacterium sp.	2	2
Klebsiella spp.	3	2	Streptococcus oralis	1	1
Citrobacter koseri	3	2	Micrococcus luteus	1	1
Candida parapsilosis	1	1	Diphteroid bacilli	1	1
Escherichia coli	1	1	Anaerobic streptococci	1	1
Pseudomonas spp.	1	1	Streptococcus species (Facklamia Languida)	1	1
Staphylococcus aureus	1	1	Streptococcus parasanguinis	1	1
Haemophilus influenzae	1	1	Granulicatella adiacens	1	1
	32	20	Saccharomyces cerevisiae	1	1
			Actinomyces sp.	1	1
			Corynebacterium striatum	1	1
			Lysinbacillus sphaericus	1	1
				60	49

#### Legend:

**n**= Number of times an organism was cultured from a test sample

BAL culture					
Likely pathogen	n	N	Likely contaminant	n	N
Klebsiella spp.	17	14	Candida spp	21	15
Escherichia coli	7	5	Enterococcus spp	2	2
Pseudomonas spp.	5	4	Yeast	2	2
Enterobacter spp.	4	3	Upper respiratory tract flora	1	1
MRSA	4	2	Streptococcus anginosus	1	1
Serratia marcesens	3	2		27	21
Staphylococcus aureus	3	3			
Pluralibacter gergoviae	2	1			
Proteus mirabilis	2	2			
Citrobacter koseri	2	2			
Raoultella sp.	1	1			
Morganella morganii	1	1			
Stenotrophomonas maltophilia	1	1			
Haemophilus influenzae	1	1			
	53	42	<u>_</u>		

### Legend:

**n**= Number of times an organism was cultured from a test sample

Likely pathogen	n	N	Likely contaminant	n	N
Serratia marcesens	10	5	Candida spp	10	6
Enterobacter spp.	7	3	Yeast	6	4
Escherichia coli	6	2	Mixed growth of Coliform & Candida	4	2
Klebsiella spp.	4	3	Respiratory commensals	2	1
Raoultella sp.	4	1	Corynebacterium sp	1	1
Pseudomonas spp.	3	2		23	14
Proteus mirabilis	2	2		<del></del>	
Staphylococcus aureus	2	2			
Haemophilus influenzae	2	1			
Aspergillus fumigatus	2	1			
Citrobacter koseri	1	1			
Acinetobacter baumanii	1	1			

#### Legend:

**n**= Number of times an organism was cultured from a test sample

Sputum culture			
Likely pathogen	n	N	Likely contamina
Pseudomonas spp.	19	11	Candida spp
Stenotrophomonas maltophilia	7	3	Enterococcus sp
Escherichia coli	6	5	Respiratory commensals
Klebsiella spp.	8	6	
Citrobacter koseri	6	5	
Enterobacter spp.	6	6	
Staphylococcus aureus	7	6	
Proteus mirabilis	2	2	
Serratia marcesens	2	2	
Burkholderia multivorans	2	1	
Haemophilus influenzae	2	1	
Pluralibacter gergoviae	1	1	
Delftia acidovorans	1	1	
Yersinia enterocolitica	1	1	
Acinetobacter baumanii	1	1	_
	71	52	

### Legend:

**n**= Number of times an organism was cultured from a test sample

	Co-infection rate within 48 hours (1000 person-days) (95% CI)
Overall	28.2 (16.7-47.7)
Excluding Nottingham University Hospitals	32.0 (17.7-57.8)

Type of tests	On admission		After admission (within 48 hours)		Overall	
	n	N	n	N	n	N
Blood culture	223	174	89	46	312	220
BAL PCR/ culture, sputum culture, tracheal culture	18	15	34	31	52	46
Urinary pneumococcal antigen	25	25	55	51	80	76
Urinary legionella antigen	36	34	55	54	91	88
Respiratory viral PCR	119	106	32	22	151	128

### Legend:

**n**= Number of tests done

**N**= Number of patients who had the test

	AMR	No AMR	Unknown	Total	Resistance n (%)
Klebsiella spp.	10	4	2	16	Co-amoxiclav, 2 (20); Cefuroxime, 7 (70); Piperacillin/Taz, 5 (50); Meropenem, 1 (10); Co-trimoxazole, 1 (10); Trimethoprim, 1 (10); Chloramphenicol, 1 (10)
Escherichia coli	11	9	0	20	Amoxicillin, 8 (72.7); Co-amoxiclav, 5 (45.5); Meropenem, 1 (9.1); Ertapenem, 1 (9.1)
Enterobacter aerogenes	3	2	0	5	Cefuroxime, 2 (40); Cefadroxil, 1 (20); Ceftazidime, 1 (20); Meropenem, 1 (20); Gentamicin, 1 (20)
Pseudomonas spp.	7	4	2	13	Ciprofloxacin, 2 (28.6); Ceftazidime, 2 (28.6); Piperacillin/Taz, 5 (71.4); Meropenem, 3 (42.9); Gentamicin, 1 (14.3); Amikacin, 1(14.3); Ticarcillin/ clavulanate, 1 (14.3)
Serratia marcescens	1	0	0	1	Piperacillin/Taz, 1 (100)
Citrobacter koseri	1	3	1	5	Piperacillin/Taz, 1(100); Meropenem, 1 (100)
Staphylococcus aureus	4	7	0	11	Flucloxacillin, 1 (25); Doxycycline, 2 (50); Clarithromycin, 3 (75); Clindamycin, 1(25)
Haemophilus influenzae	3	0	1	4	Amoxicillin, 3 (100); Co-amoxiclav, 3 (100); Cefuroxime, 2 (66.7); Doxycycline, 1 (33.3)
Acinetobacter baumanii	1	1	0	2	Ceftazidime, 1 (100)
Burkholderia multivorans	1	0	0	1	Meropenem, 1 (100); Ceftolozane/Tazobactam, 1 (100)
Enterococcus spp.	1	3	4	8	Amoxicillin, 1 (100); Gentamicin, 1 (100)
Morganella morganii	1	0	0	1	Cefuroxime, 1 (100); Piperacillin/Taz, 1 (100)
Raoultella sp.	1	0	0	1	Piperacillin/Taz, 1 (100)
Total	45	33	10	88	

#### Legend

- AMR is defined as resistance reported to one or more antimicrobial agents tested (excluding intrinsic resistances).(2) Information in this table is based on the antimicrobial patterns released by individual sites for clinicians, other resistance may have been present but not reported
- Piperacillin/Taz= Piperacillin/Tazobactam
- Co-pathogens (with AMR) breakdown by species
  - o Klebsiella spp.: Klebsiella pneumoniae (2), Klebsiella aerogenes (7), Klebsiella voriicola (1)
  - o Pseudomonas spp.: Pseudomonas sp (1), Pseudomonas aeruginosa (6)
  - o Enterococcus spp.: Enterococcus faecium (1)